

## CHANGES IN THE ANNUAL CYCLE AND SEASONAL CHARACTERISTICS OF PRECIPITATION IN THE DANUBE RIVER LOWER BASIN

Ancuta MANEA<sup>1</sup>, Nina. NIKOLOVA<sup>2</sup>,  
Constanta BORONEANT<sup>1</sup>, Alina ORZAN<sup>1</sup>

<sup>1</sup>National Meteorological Administration, Bucharest, Romania

<sup>2</sup>St. Kliment Ohridsky University of Sofia, Bulgaria

[e-mail:ancuta.manea@meteoromania.ro](mailto:ancuta.manea@meteoromania.ro)

( Received October 2010, final version November 2011)

**Abstract:** Precipitation is one of the main climatic elements which determines the availability of water resources and has considerable impact on different aspects of human activity. The present analysis is based on monthly precipitation totals from 22 meteorological stations situated in the Danube River lower basin both in Romania and Bulgaria. These stations have been divided into three groups by their geographical distribution and the peculiarities of the study area (western, central and eastern). The aim of this paper is to provide detailed information about the characteristics of annual and seasonal variability of precipitation in this area, focusing on annual cycle and linear trends. The investigated period is 1961-2007. To reveal the characteristics of temporal variability of precipitation, two 30-year reference periods have been considered: 1961-1990 and 1978 – 2007. In general, for the entire analyzed area and complete period 1961-2007, the annual cycle of precipitation showed June as the wettest month and February and October as the driest months. However, as regards the details on each subgroup, some changes of the annual cycle occurred between subgroups and the two reference periods. For instance, in the western subgroup, the wettest month on record is June for 1961-1990 and May for 1978 – 2007. In the central subgroup, it is still June for 1961-1990, but it is July for 1978-2007. Regarding the linear trend over 1961-2007, a decrease for the winter season was identified, which is statistically significant only at six stations. For other seasons, no statistically significant trends were found.

### 1. INTRODUCTION

Precipitation is one of the main climatic elements which determines the availability of water resources and has considerable impact on different aspects of human activity. Knowledge of seasonal precipitation characteristics can bring further understanding of peculiarities of precipitation regime in a region of interest. The scientific works (Hulme et al., 1998) show that there are

opposite trends in precipitation variability in northern Europe (wetting) and southern Europe (drying).

Seasonal precipitation in Romania has been investigated in connection with the large-scale circulation patterns (Busuioc and von Storch, 1996; Rimbu et al, 2001; Tomozeiu et al, 2005) It was shown that some types of atmospheric circulation, large scale mechanisms such as the North Atlantic Oscillation and El Niño-Southern Oscillation can be

responsible for the anomalies in precipitation regime over the Romanian territory and, implicitly, can impact on the Danube River flow in its lower basin (Rimbu et al. 2002; 2004). Records of temperature and precipitation anomalies in Romania during the recent years have been analyzed in the context of corresponding extremes since the beginning of the 20th century (Busuioc et al. , 2007) and the results show that the frequency and the magnitude of extreme events have increased both for temperature and precipitation in the last decades (Busuioc et al. 2003; 2007).

The Balkan Peninsula is characterized by general decreasing trends of precipitation beginning with the 1980s (Alexandrov, 2004). As it is an important issue, variability in Bulgaria's precipitation regime has been studied by many authors (Topliiski, 2002, 2005, Zlatunova and Penkov, 2002, Vekilska and Rathcev, 2000). There are several publications dedicated to precipitation variability across the Danubian Plain (Koleva, 1995; Nikolova and Vassilev, 2006; Petkova et al., 2008).

Though many publications on trends and seasonality of precipitation in Europe put the Balkan region and the south-eastern Europe in the context of global changes (Zveryaev, 2004), an in depth statistical analysis is needed in order to get more insight into the precipitation regime in our region in the context of contemporary climate change. The aim of this paper is to provide detailed information about the characteristics of annual and seasonal variability of precipitation in the Danube River lower basin, focusing on annual cycle and linear trends.

## 2. DATA AND METHODS

The present analysis is based on monthly precipitation totals from 22 meteorological stations situated in the Danube River lower basin both in Romania and Bulgaria. Data for Romanian stations are provided by the National Meteorological Administration, Romania. Data for Bulgarian stations have been compiled from the meteorological yearbooks published by the National Institute of Meteorology and Hydrology, Sofia, Bulgaria, from the Statistical Yearbook of the National Statistical Institute in Bulgaria and from the bulletins issued by Bulgaria's Executive Environmental Agency. The geographical coordinates of the stations are presented in Table 1. The stations have been divided into three groups according to their geographical distribution and the peculiarities of the study area as follows: 1) western part of the domain which includes the stations: Tr. Severin, Calafat, Băilești, Bechet, Caracal, Tr. Măgurele, Vidin, Vratsa, Lom, Oryahovo, Kneja and Pleven; 2) central part of the domain including the following stations: Alexandria, Giurgiu, Obratzov Chiflik, Razgrad, Bucuresti Băneasa and Călărași and 3) north-eastern part of the domain including the stations: Sulina, Tulcea, Galați and Hârșova.

The investigated period is 1961-2007. To reveal the characteristics of temporal precipitation variability, two reference periods have been considered: 1961-1990, which is named *baseline climate* by the World Meteorological Organization and the last 30 years (1978 – 2007), referred to as *recent climate*.

CHANGES IN THE ANNUAL CYCLE AND SEASONAL CHARACTERISTICS OF  
PRECIPITATION IN THE DANUBE RIVER LOWER BASIN

**Table 1.** Geographical coordinates of the meteorological stations considered in the present study.

	Station name	Lat;	Lon	Altitude	Country
1.	Tr. Severin	44° 38'	22° 38'	77	RO
2.	Calafat	43° 59'	22° 57'	61	RO
3.	Baileşti	44° 01'	23° 20'	57	RO
4.	Bechet	43° 47'	23° 57'	36	RO
5.	Caracal	44° 06'	24° 22'	106	RO
6.	Tr. Măgurele	43° 45'	24° 53'	31	RO
7.	Vidin	43° 39'	22° 53'	37	BG
8.	Vratsa	43° 12'	23° 32'	548	BG
9.	Lom	43° 49'	23° 14'	21	BG
10.	Oryahovo	43° 44'	23° 58'	31	BG
11.	Kneja	43° 30'	24° 05'	120	BG
12.	Pleven	43° 25'	24° 35'	133	BG
13.	Alexandria	43° 59'	25° 21'	75	RO
14.	Giurgiu	43° 53'	25° 57'	24	RO
15.	Obraztsov chiflik	43° 48'	26° 02'	153	BG
16.	Razgrad	43° 32'	26° 31'	183	BG
17.	Bucharest-Băneasa	44° 30'	26° 08'	90	RO
18.	Călăraşi	44° 12'	27° 21'	19	RO
19.	Sulina	45° 10'	29° 44'	3	RO
20.	Tulcea	45° 11'	28° 49'	4	RO
21.	Galaţi	45° 29'	28° 02'	69	RO
22.	Hârşova	44° 41'	27° 57'	38	RO

To obtain the most reliable results the quality of initial data has been verified. The data have been checked for missing values and have been tested for homogeneity. There are not missing values in the investigated time series.

A homogeneous time series is defined as one in which trend and variability are not influenced by artificial factors such as relocation or environmental changes around the station, changes of instruments or of the observation methods (Keiser and Griffiths, 1997; Sneyers, 1975).

The AnClim software (<http://www.climahom.eu/AnClim.html>) has been used to test the homogeneity of monthly precipitation with the Alexandersson test (Alexandersson and Moberg, 1997). Non-homogeneities (referred to afterwards as breakpoints) have been detected mainly after the 80s.

For the Bulgarian stations this may be due to the different data sources.

The distribution of breakpoints along the months of a year (Figure 1) shows that the maximum number of stations (9) have been detected in April. At 5 stations break points have been detected in June and at 4 stations in March, May and July, respectively.

All the series of monthly precipitation totals where breakpoints were detected have been adjusted with the AnClim software. After the homogenization of monthly precipitation totals the seasonal and annual totals have been calculated. The seasons are determined as follows: winter – December, January, February; spring – March, April, May; summer – June, July, August and Autumn – September, October, November.

In order to determine the trend of series of seasonal and annual precipitation totals, linear regressions have been used.

precipitation amount in summer months can be associated with convective precipitation that is usually observed in this season. In general, the monthly




Fig. 1. Number of stations with breakpoints during a year over 1961-2007.

### 3. RESULTS

#### 3.1. Annual cycle


The analysis of the results for the Romanian and Bulgarian meteorological stations considered together for the entire domain revealed that the highest monthly average in the annual cycle of precipitation was, in general, recorded in June but at some stations the rainiest month is May or July, with some differences between the two periods. During the 1961-1990 time period the analysis shows an important concentration of the highest monthly averages in June. The 1978 – 2007 period does not display the same stability feature for June. The maximum values are distributed almost evenly over May, June and July. The monthly minimum precipitation is mainly recorded in February and October. (Figure 2, an example for some stations). The highest

precipitation amounts over 1978-2007 are lower than those for the reference period 1961-1990, except for the autumn months (September, October, November).


The distribution of the highest and lowest monthly averages was analyzed for each subgroup of stations in order to determine the general and regional characteristics of the wettest and driest months.

The analysis reveals different characteristics of the annual cycle for each subgroup and some differences between the two sub-periods. During 1961-1990, the wettest month is June for the western and central subgroups, while for the eastern one, the wettest month is May. This characteristic changed in the second period (1971-2007), since the wettest month on record was as follows: May-western subgroup, July-central subgroup and June- eastern subgroup (Figure 3).

## CHANGES IN THE ANNUAL CYCLE AND SEASONAL CHARACTERISTICS OF PRECIPITATION IN THE DANUBE RIVER LOWER BASIN



**Fig. 2.** Annual cycle of monthly precipitation totals for the group of stations located in the central part of the Danube River lower basin. Comparison between the periods 1961-1990 and 1978-2007.



**Fig. 3.** Distribution of the highest monthly precipitation averages for 1961-1990 (left column) and 1978 – 2007 (right column) with time periods for each subgroup.

Regarding the driest month over the 1961-1990 period, records showed it to be September –western subgroup, October-central subgroup and an even distribution between March and October-eastern subgroup. Over 1971-2007, the driest month on records is February for each subgroup (Figure 4).

summer precipitation has the main contribution in the annual precipitation totals (Clima Romaniei, 2008). The summer precipitation amount accounts for about 30% of the annual precipitation (Figure 5). The spring precipitation total contributes with about 25% to the annual precipitation and the winter precipitation




Fig. 4. Distribution of the lowest monthly precipitation averages over 1961-1990 and 1978 – 2007 for each subgroup.




Fig. 5. Seasonal distribution of precipitation totals (% of annual values) for each subgroup.

Comparisons between seasonal and annual precipitation amounts over the entire period 1961-2007, show that

total accounts for about 20% and even less of the annual total.

CHANGES IN THE ANNUAL CYCLE AND SEASONAL CHARACTERISTICS OF  
PRECIPITATION IN THE DANUBE RIVER LOWER BASIN

### 3.2. Trends

An analysis of the linear trend over 1961-2007 pointed out some characteristics of the precipitation regime for our region of interest (Table 2).

A decreasing trend in winter precipitation is statistically significant for some of the stations from the western and eastern groups (Table 2).

In spring and summer the trend is not statistically significant (not shown). For autumn (Table 2), a slightly positive trend has been identified, mainly due to the increase occurred after 1978, but statistically significant only for two stations in the central subgroup.

At most of the investigated stations a decreasing trend (but not statistically significant) in annual precipitation totals

has been detected over 1961-2007. Only six stations in the central subgroup (Alexandria, Giurgiu, Bucharest-Baneasa, Obratzsov chiflik, Razgrad and Calarasi) show an increasing (not statistically significant) trend (Table 2).

The results presented above show that decadal variability is the most important characteristic of the precipitation amount in the analyzed area. This result is in agreement with that presented by Busuioc et al. (2010) in a study covering the entire Romania.

### 4. CONCLUSIONS

In general, the annual cycle of precipitation, showed that June is (as) the wettest month and February and October

**Table 2.** Linear trend of winter, autumn and annual precipitation (mm/10 years) over 1961-2007. The values marked with asterisk are statistically significant at significant level  $p < 0.05$ .

	winter	autumn	annual
Tr. Severin	-13.5	-3.1	-24.1
Calafat	-8.4	7.8	2.5
Bailești	-12.6*	6.6	-2.7
Bechet	-31.8*	1.4	-4.6
Caracal	-14.4*	6.5	-15.6
Tr. Măgurele	-12.9*	6.4	-13.9
Vidin	-5.9	3.6	-9.3
Vratsa	-10.1*	1.6	-12.3
Lom	-9.5	8.5	-0.9
Oryahovo	-9.2	-1.6	-12.8
Kneja	-6.9	1.0	-8.5
Pleven	-12.0*	4.5	-6.7
Alexandria	-5.6	12.2	11.0
Giurgiu	-4.8	13.5	5.5
Obratzsov chiflik	-5.9	13.0	6.2
Razgrad	-4.5	16.6*	7.0
Bucharest - Băneasa	-7.5	15.8	6.1
Călărași	24.5	15.6*	11.1
Sulina	-8.3*	9.5	5.2
Tulcea	-13.6*	2.1	-6.9
Galați	-10.5*	8.3	5.4
Hârșova	-14.0*	-2.1	-22.0

the driest ones, with some changes for the two analysed sub-periods and the three regional subgroups. The analysis over the entire considered area showed some particularities for each 30-year period:

- the wettest month is very well represented by June for the 1961-1990 time period;
- the driest month reaches more than 80% of the number of cases in February for the 1978 – 2007 time period.

However, a detailed analysis over the three subgroups revealed that, during 1961-1990, the wettest month is June for the western and central subgroups, while for the eastern one, the wettest month is May. For 1971-2007, the wettest month is recorded in May-western subgroup, July-central subgroup and June-eastern subgroup.

Regarding the driest month, for the 1961-1990 period, this is recorded in

September –western subgroup, October-central subgroup and evenly distributed between March and October-eastern subgroup. For the period 1971-2007, the driest month is recorded in February for all subgroups.

As regards the seasonal characteristics, one of the most important findings is the identification of a decreasing trend in winter precipitation but statistically significant only at six stations. For other seasons, no statistically significant trends were found. This result shows that decadal variability is the most important characteristic of the precipitation amount in the analyzed area. This result is in agreement with that presented by Busuioc et al. (2010) in a study which covers the entire Romania.

**Acknowledgements:** This study was made possible by funds taken from a collaboration project between Romania and Bulgaria (Contract nr. 100/05.09.2008). Two anonymous reviewers are acknowledged for their useful comments.

CHANGES IN THE ANNUAL CYCLE AND SEASONAL CHARACTERISTICS OF  
PRECIPITATION IN THE DANUBE RIVER LOWER BASIN

**REFERENCES**

- Alexandersson, H. Moberg, A. (1997).  
*Homogenization of Swedish temperature data. 1. Homogeneity test for linear trends.* Int. J. Climatol., 17: 25-34.
- Alexandrov, V., 2004.  
Climate variability and change and related drought on Balkan Peninsula. Proceedings of the Conference on Water Observation and Information System for Decision Support (BALWOIS) Ohrid, Macedonia, 25-29 May 2004 (CD version)
- Busuioc, A., Caian, M., Cheval, S., Bojariu, R., Boroneant, C., Baciu, M., Dumitrescu, A., 2010: *Variabilitatea si schimbarea climei in Romania*, **Ed. Pro Universitaria**, ISBN: 978-973-129-549-7, Bucuresti, 226 pp.
- Busuioc A., C. Boroneant, D. Matei, 2003:  
Analiza variabilitatii frecventei evenimentelor pluviometrice extreme în Romania în „Analele Universității Ovidius, Seria: Geografie”, vol. 1, 2003, 43-50.
- Busuioc, A., Al. Dumitrescu, E. Soare, A. Orzan, (2007):  
*Summer anomalies in 2007 in the context of extremely hot and dry summers in Romania*, Romanian Journal of Meteorology, Vol. 1. No 1-2, 1-17.
- Busuioc, A., H. Von Storch, 1996:  
"Changes in the winter precipitation in Romania and its relation to the large scale circulation", *Tellus*, 538-552.
- Hulme, M., T.J. Osborn, & T.C. Johns, 1998:  
Precipitation sensitivity to global warming: comparison of observations with HadCM2 simulations. *Geophysical Research Letters*, 25, 3379–3382
- Keiser, D. T., J. F. Griffiths. 1997.  
Problems associated with homogeneity testing in climate variation studies: a case study of temperature in the northern Great Plains, USA. *Int. J. of Climatology*, Vol. 17, Issue 5 , Pages 459 – 563.
- Koleva, E. 1995.  
Drought in the lower Danube basin. *Drought network news*. Vol. 1, N 1, Feb. Nebraska, USA, 6-7
- Nikolova, N., S. Vasilev. 2006.  
*Variability of Summer-Time Precipitation in Danube Plain, Bulgaria*. Collection of papers, Geographical Institute “Jovan Cvijic” Serbian Academy of Sciences, N 54.

- Petkova, N., E. Koleva, V. Alexandrov. 2008.  
 Winter Climate Variability at Danube Plane, North Bulgaria. XXIVth Conference of the Danubian Countries on the Hydrological Forecasting and Hydrological Bases of Water Management. Bled, Slovenia. 2 – 4 June 2008. Conference e-papers. ISBN 978-961-91090-2-1
- Rimbu N., C. Boroneant, C. Buta, M. Dima, (2002):  
*Decadal variability of the Danube river streamflow in the lower basin and its relation with the North Atlantic Oscillation.* International Journal of Climatology, 22, 1169-1179.
- Rimbu N., H. Le Treut, S. Janicot, C. Boroneant, and C. Laurent, (2001):  
*Decadal precipitation variability over Europe and its relation with surface atmospheric circulation and sea surface temperature,* Quart. J. Royal Met. Soc., vol. 127, no. 572, part B, 315-329.
- Rimbu, N., Dima, M. Lohmann, G. and Stefan, S. (2004)  
 Impacts of the North Atlantic oscillation and the El Nino-southern oscillation on Danube river flow variability. Geophys. Res. Lett., 31, L23203, doi:10.1029/2004GL020559.
- Sneyers, R., 1975:  
 Sur l'analyse statistique des series d'observation, *WMO Note Technique*, no. 143, 189 pp.
- Stefan, S., Ghioca M., Rimbu N., C. Boroneant, (2004):  
*Study of meteorological and hydrological drought in southern Romania from observational data.* International Journal of Climatology, 24, 871-881
- Tomozeiu, R., Stefan, S. and Busuioc, A., 2005:  
 Spatial and temporal variability of the winter precipitation in Romania in connection with the large-scale circulation patterns, *Theoretical and Applied Climatology* DOI 10.1007/s00704-004-0082-3.
- Topliiski, D., (2002):  
 Climatic changes in Bulgaria in the period 1901 – 1990. Proceedings of the International Scientific Conference in Memory of Prof. Dimitar Yaranov, Varna, Bulgaria, 98-106. (in Bulgarian).
- Topliiski, D., (2005).  
*Chronological Structure of Climate in Bulgaria during 20 Century.* Dissertation for Dr Sc. (in Bulgarian)
- Vekilska, B., G. Rathcev, (2000).  
*Current Changes in the Precipitation in Bulgaria.* Sofia University Year Book, Vol. 90, Geography., 31-37.
- Zlatunova, D. , I. Penkov. (2002).  
 River runoff and climate change in Bulgaria – Proceedings of the International Conference “Preventing and Fighting Hydrological Disasters”, 2002, Timisoara, Romania. 413-416.
- Zveryaev, I. I. (2004),  
*Seasonality in precipitation variability over Europe,* J. Geophys. Res., 109, D05103, doi:10.1029/2003JD003668.